Table 3.—Maximum free air wind velocities, (m. p. s.), for different sections of the United States based on pilot-balloon observations during October 1941

Section	Surface to 2,500 meters (m. s. l.)					Between 2,500 and 5,000 meters (m. s. l.)					Above 5,000 meters (m. s. l.)				
	Maximum ve- locity	Direc- tion	Altitude (m.) m. s. l.	Date	Station	Maximum ve-	Direc- tion	Altitude (m.) m. s. l.	Date	Station	Maximum ve- locity	Direc- tion	Altitude (m.) m. s. l.	Date	Station
Northeast 1	42. 6 36. 2 26. 3 39. 4 36. 6 34. 0 36. 0 33. 4 29. 9	{WSW SW SW WSW SW WSW NNW	1,740 2,420 1,650 2,100 2,500 1,970 1,900 430	5 8 27 27 7 5 17 3 2 5	Sacramento, Calif	28. 2 63. 2 45. 0 38. 4 37. 6 51. 6	SSW SW WSW NNE	4, 400 3, 940 3, 170 4, 970	23 10 10 7 4 26 10 2 13 2	Kylertown, Pa Huntington, W. Va. Spartanburg, S. C Muskegon, Mich Wichita, Kans Oklahoma City, Okla. Butte, Mont Reno, Nev Winslow, Ariz Bakersfield, Calif	70.0 55.2 74.5 73.0	WNW WNW WNW SW WNW	8,820 11,570 20,500 11,560 9,170 13,255 10,220 11,480 12,320	17 28 20 10 9 31 2 25 8	Albany, N. Y. Greensboro, N. C. Miami, Fla. Minneapolis, Minn Omaha, Nebr. San Antonio, Tex. Great Falls, Mont. Reno, Nev. Las Vegas, Nev.

## WEATHER ON THE NORTH ATLANTIC OCEAN

By H. C. HUNTER

Atmospheric pressure.—The pressure during October 1941, averaged above normal over large portions of the North Atlantic, especially near the coasts of the South Atlantic States, Cuba, and the Bahamas. On the other hand it averaged below normal near the Maritime Provinces, Newfoundland, and Labrador.

The extremes of pressure noted in vessel reports at hand were 1,036.2 and 991.5 millibars (30.60 and 29.28 inches). The high reading was noted during the forenoon of the 30th, about 50 miles from Montauk Point, in a south-southwest direction. Table 1 shows that the pressure was slightly higher at Halifax, Nova Scotia, that day. The low mark was noted near southeastern Newfoundland during the early afternoon of the 20th. During other portions of the month both Belle Isle and Halifax recorded lower pressures.

In low latitudes readings slightly below 1,000 millibars (29.53 inches) were noted by two vessels near 29° N., 75° W., on the 10th, when within the area affected by a storm of tropical origin. However, this storm, when crossing the Bahama Islands several days earlier, as indicated in an article on page 303 of this Review, resulted in a pressure of 964.4 millibars (28.48 inches) on Cat Island.

Table 1.—Averages, departures, and extremes of atmospheric pressure (sea level) at selected stations for the North Atlantic Ocean and its shores, October 1941

		··· <del>·</del>				
Station	Average pressure	Depar- ture from normal	Highest	Date	Lowest	Date
Horta, Azores Belle Isle, Newfoundland Halifax, Nova Scotia Nantucket Hatteras Turks Island <sup>1</sup> Key West New Orleans	Millibars 1, 020. 0 1, 007. 9 1, 014. 8 1, 018. 0 1, 020. 3 1, 016. 0 1, 016. 3 1, 017. 3	Millibars +0.4 -3.3 -2.5 +.4 +2.3 +1.8 +2.4 +.4	Millibars 1, 031 1, 029 1, 038 1, 036 1, 030 1, 019 1, 022 1, 024	13 30 30 30 29 18 14	Millibars 1, 009 986 989 1, 000 1, 009 1, 012 1, 011 1, 010	7 12 11 10 10 16 6

<sup>1</sup> For 27 days.

Indiana, Illinois, Iowa, Nebraska, Kansas, and Missouri.
 Mississippi, Arkansas, Louisiana, Oklahoma, Texas (except El Paso), and Western

\*\* Montana, Idaho, Washington, and Oregon.

\*\* Wyoming, Colorado, Utah, Northern Nevada, and Northern California.

\*\* Southern California, Southern Nevada, Arizona, New Mexico, and extreme West

Cyclones and gales.—The vessel reports available indicate a comparatively quiet October. In middle latitudes two strong gales and several fresh gales occurred. The period centering on the 20th appears to have been the most eventful, and from the 8th to 11th likewise was somewhat unquiet.

The opening week and the period from 22d to 26th seem to have been remarkably undisturbed from the 30th parallel northward.

Tropical disturbances.—In another portion of this issue an account is presented of two disturbances of tropical origin. The earlier of these cyclones was noted to northward of the Virgin Islands on the 3d; it moved westward across the Bahamas and extreme southern Florida, turned northward over the Gulf and traversed parts of western Florida, southern Georgia and South Carolina, then moved southeastward over the Atlantic to near latitude 30°, thence continued mainly eastward till beyond the field of

This storm was of comparatively small diameter during much of its course, and was for awhile remarkably intense, especially over the Bahamas. A few lives are reported lost in the Bahamas and some in northwestern Florida. No vessel report that relates to this storm indicates a higher wind than a strong gale (force 9).

The later LOW was of short path and little moment. Its track was approximately the middle portion of the track of the earlier storm, and the time was about 13 days later. The greatest wind force noted in vessel reports as a result of this disturbance was a moderate gale (force 7).

Line squalls.—About 150 miles to eastward of the extreme southeastern coast of Maryland line squalls were noted from a vessel which furnishes this report:

October 1, 9:30 p. m., 75th meridian time (equivalent to October 2, 2:30 a. m., Greenwich mean time), in latitude 38°04′ N., longitude 73°00′ W., barometer 30.16 inches (1021.3 millibars), having risen 0.03 inch during 1½ hours preceding, temperature of air 72°, of water 70°; three line squalls, moving southeastward, ranging from about 15 to 20 miles in length, passed at intervals of about 5 to 8 minutes. The clouds appeared in a very compact mass and in an almost straight line. During this time the wind shifted from southalmost straight line. During this time the wind shifted from southwest, Beaufort force 3, to northwest, same force; within the succeeding 3½ hours the wind turned partly back, becoming west, force 2. Other clouds at the time of the squalls were alto-cumulus, about two-tenths of sky being covered, apart from the line squall masses which covered a third of the overhead as they passed. Sea small southwest. Ship's course north-northeast, speed 14 knots.

Maine, Vermont, New Hampshire, Massachusetts, Rhode Island, Connecticut, New York, New Jersey, Pennsylvania and Northern Ohio.
 Delaware, Maryland, Virginia, West Virginia, Southern Ohio, Kentucky, Eastern Tennessee and North Carolina.
 South Carolina, Georgia, Florida, and Alabama.
 Michigan, Wisconsin, Minnesota, North Dakota, and South Dakota.

Note.—All data based on available observations, departures compiled from best available normals related to times of observation, except Hatteras, Key West, Nantucket, and New Orleans, which are 24-hour corrected means.

Waterspouts near the Equator.—Approximately 700 miles to northward of Cape St. Roque (in 5°10′ N., 33°04′ W.), on October 14, about 7:50 a.m., local time, with barometer 1,013.9 millibars (29.94 inches), waterspouts were observed, as reported below:

Wind fitful and light, shifting continuously from north to east to southwest and back. Partly cloudy and light rain squalls. Cloud movements shifty and in various directions.

One mass of clouds, apparently at a lower level than the surrounding cloud formations, drifting from a northeasterly direction, moved into what seemed to be a stationary mass of clouds, almost immediately forming a single spout close enough to our vessel that the boiling motion on the surface of the sea could be observed. As this spout seemed to gain in violence, and as the column darkened, it

appeared to move in a northwesterly direction.

Perhaps 5 minutes later the column of the spout began to lighten

and fade.

At this time two more spouts formed in the same cloud group further to the southeast. These two columns, one of which was very twisted, seemed to start fading almost as soon as they were well formed. These two lasted perhaps 5 minutes.

The first spout was in formation a little better than 10 minutes when it started fading. The column of the spout seemed to part in the center, the lower half dropping into the sea, and the upper half receding upward into the cloud. Some time after the lower half had dropped into the sea, however, and while the upper half was still fading upward, a motion was still visible on the surface of the sea.

Past weather was with light rain squalls, and the prevailing cloud formation was cumulonimbus.

Fog.—There was but little fog reported this month. The square 40° to 45° N., 65° to 70° W., had fog on 3 days, and a very few other ocean areas are indicated as having fog on 1 or 2 days. No fog was reported from any position to eastward of the 55th meridian, or to southward of the 35th parallel.

Even where most fog was noted, near the coasts of eastern New England and of Nova Scotia, the occurrence was apparently less than the normal during the month of October.

## OCEAN GALES AND STORMS, OCTOBER 1941

CCEAN GABES AND STORMS, OCTOBER 1941											
Vessel		at time of arometer	Gale began Octo-	Time of lowest barometer,	Gale ended, Octo-	Lowest ba- rometer	Direction of wind when gale began	Direction and force of wind at time of lowest	Direction of wind when gale ended	Direction and highest force of wind	Shifts of wind near time of lowest barometer
	Latitude	Longitude	ber		ber			barometer	<b>G</b>	wind	
NORTH ATLANTIC OCEAN	. ,	0 /				Millibars			_		
A vessel	24 12N. 25 42N.	69 18W. 77 12W.	4 5	1p, 4 7p, 5	4 5	1, 007. 8 1, 005. 3	ENE	ENE, 7 ENE, 8	SESE.	SE, 8 ENE, 8	ENE-E. NE-E.
Do	27 36N. 37 06N.	82 42W. 69 24W.	6 8	4p, 6 2p, 8	6 8	1, 005. 8 1, 010. 8	NE	SE, 9 W, 8		SE, 9 W, 8	NE-SE.
Do Do	32 10N. 38 06N.	77 30W. 58 18W.	8	5p, 8	8	1, 002. 7 1, 004. 4	SSW	ESE, 9 SW, 9 NW, 6	Esw	LESE.9	S-E. SW-NNW.
Do Do	29 12N. 27 54N.	74 42W. 79 42W.	10 11	2p, 10	10 11	999. 3 1. 012. 9	ESE	IN.3	E	SW, 9 ESE, 8 N, 8	ESE-NNE-W.
Do Do	40 42N. 38 30N.	68 06W. 60 06W.	10 11	10p, 10 8a, 11	11 11	1, 002. 7 1, 002. 0	NW	WNW,7 SSW,8 SW,7	NW	NW, 8 SW. 8	ssw-nw.
Do	29 00N. 39 06N.	70 00W. 47 48W.	11 20	8a, 11 9a. 20	12 21	1, 003. 1 997. 0	SW	SW, 7	NNE	N, 8 NW, 8	SW-W. SSE-SSW-NNW.
Do Do	47 30N. 39 06N.	52 40W. 60 42W.	19 28	1p, 20 2a, 29	20 29	991. 5 1, 015. 9	SSE	N. 9 SSW. 8	N	N. 9. SSW. 8	NNE-N. S-SSW.
NORTH PACIFIC	05 0011.	00 12 17 .	2.5	24, 20	20	1,010.0		55 11, 6		0011,011111	B SBW.
OCEAN A vessel	42 36N.	169 42E.	1	5p, 30 1	1	1, 019, 6	NNW	NNW. 6	N	N. 8	•
Do	38 24N.	124 00W.	į	4p, 1	2 2	1, 015. 9	NW	NW, 7	NW	NW, 8 SW, 8	, ozar
Do Do	53 54N. 37 36N.	156 12W. 124 54W.	1	8a, 1	2	994.6 1,015.6	SW. WNW	SW, 6 NW, 8	WSW	NW. 8	S-SW. None.
Do Do	57 30N. 48 00N.	151 00E. 168 24W.	2 3	3a, 4	3 4	997. 0 1, 013. 2	WNWs	NW, 5 S, 7	WNW	WNW, 8 S, 8	WNW-NW.
Do	49 18N. 48 19N.	132 48W. 124 49W.	8 9	9a, 9 12m, 9	9	999. 3 1, 003. 4	WSE	SW, 7 SE, 8	WSW	W, 8 SE, 8	E-SW-SSW.
Do	30 36N. 24 21N.	149 36E. 156 20E.	10 10	4a, 11	11 13	1, 018. 0 993. 3	N	NE, 8 NE, 10	NE	NE, 8 NE, 10	None.
Do Do	31 00N.	154 00E.	12	6a, 13	13	1, 009. 1	ENE	ENE. 7	ESE	ENE. 8	ENE-ESE.
Do Do		148 54W. 94 06W.	12 12	4p, 12 4p, 12	13 13	977.0 1,008.8	NE	N, 10 NNW, 3 S, 10	SWENE	NNE, 11 ENE, 8	NNE-WNW. W-N.
Do Do	32 48N. 51 18N.	158 00E. 138 36W.	12 14	6p, 14	15 15	1, 003. 1 985. 4	NESE.	S, 10 SW. 9	SW	8, 10 8W, 9	SSE-S. SE-W.
Do Do	2 49 29N. 54 36N.	156 56W. 135 24W.	14 15	10p, 14 9p, 15	15 16	981. 4 997. 6	SSE	SW, 8 SE, 8	wsw	WSW, 9 SE, 8	sw-wsw.
Do	47 42N.	151 12W.	15	12p, 15	18	975. 3	SE	8W, 10	WNW	SW, 10	a waw
Do Do	42 05N. 58 36N.	178 00W. 138 00W.	15 16	3a, 16 9p, 16	16	986. 8 974. 9	8	SW, 11 E, 8	NNW	SW, 11 E, 8	s-wsw.
Do	53 00N. 58 06N.	148 18W. 160 54W.	16 17	11p, 16 7a, 17	18 17	948. 2 985. 1	SE	SŚW, 10 NW, 8	W NW	SW, 10 NW, 8	S-8W. NNW-NW.
Do Do	<sup>2</sup> 51 38N. 56 18N.	138 17W. 145 06W.	15 16	4p, 17 9a, 17	19 18	988. 2 957. 3	SW	8SW, 8 ESE, 8	WSW WNW	8. 9.	S-SW. SSE-E.
Do	43 42N.	133 48W.	17	9a, 17	18	1,009.1	w	W. 4	WNW	NW, 10 WNW, 8 NW, 9	8-W.
Do	54 18N. 40 00N.	155 18W. 150 12W.	19 20	2a, 19 2p, 20	19 20	986. 1 1. 004. 4	SSE	SW, 5 SSE, 8	NW	SSE. 8	8-W.
Do	26 51N. 33 29N.	149 06W. 158 58E.	23 24	1p, 23 2a, 25	23 25	1, 011. 2 1, 009, 1	NW N	N, 10 N. 8	N	N, 10 N, 9	VarNNNW. None.
Do	40 18N. 39 30N.	138 00W. 155 12W.	24 25	6a, 25 4p, 26	25 26	984, 4 1, 005, 1	8W	8W, 8 8, 8.	SW SSW	SW, 8 S, 9.	
Do	99 9014.	100 12 17	20	7p, 20	20	1,005.1	D	D, 0	00 77	D, 8	İ

<sup>1</sup> September.

2 Position approximate.

<sup>1</sup> Barometer uncorrected.

## WEATHER ON THE NORTH PACIFIC OCEAN By WILLIS E. HURD

Atmospheric pressure.—Over the greater part of the ocean from which readings are available for October 1941, the mean pressure was practically normal. The region of greatest departure was in the Gulf of Alaska, where considerable cyclonic activity occurred. At Juneau, with a mean barometer of 1,007.1 millibars (29.74 inches), the pressure was 4.4 millibars (.13 inch) below the normal of

the month. The lowest barometer reading reported for October was 948.2 millibars (28.00 inches) read on a ship in the central Gulf of Alaska on the 16th.

There was much anticyclonic activity on the east-central part of the ocean, and the average HIGH extended from the Washington coast southwestward across Midway Island.

In southwestern waters, the island of Guam had an unusually low mean barometer for a tropical station in October. The average was 1,007.2 millibars (29.74)